

## **KUESIONER PENELITIAN**

### **A. DATA RESPONDEN**

Nama Responden :

Usia Responden :

Jenis Kelamin :

Pendidikan Terakhir :

### **B. PETUNJUK PENGISIAN PESPONDEN**

Berikut tanda checklist (√) pada kolom jawaban yang telah disediakan dibawah ini sesuai pilihan anda.

SS	: Sangat Setuju	Nilai 5
S	: Setuju	Nilai 4
KS	: Kurang Setuju	Nilai 3
TS	: Tidak Setuju	Nilai 2
STS	: Sangat Tidak Setuju	Nilai 1

### Kepemimpinan (X1)

No	Pernyataan	SS	S	KS	TS	STS
1	Pemimpin di tempat saya mengajar selalu memberikan perhatian dan motivasi kepada guru-guru untuk selalu giat dalam mengajar.					
2	Pemimpin di tempat saya mengajar memiliki strategi yang jelas dan realistis (masuk akal) dalam setiap agenda kerja yan dibuat.					
3	Pemimpin di tempat saya mengajar mampu menghargai setiap perbedaan pendapat yang diajukan oleh para guru.					
4	Saya memiliki kepercayaan kepada pemimpin saya, bahwa pemimpin mampu memberikan perubahan dan kemajuan untuk sekolah.					
5	Pemimpin di tempat saya mengajar selalu peduli terhadap permasalahan yang dihadapi oleh para guru dan memberikan arahan yang tepat.					

### Motivasi (X2)

No	Pernyataan	SS	S	KS	TS	STS
1	Pemimpin di tempat saya mengajar menerapkan pola bahwa setiap individu mampu untuk menjadi pemimpin.					
2	Saya bertanggung jawab atas suatu tugas atau pekerjaan yang saya kerjakan.					
3	Pemimpin di tempat saya mengajarsenantiasa mengarahkan untuk memperluas pergaulan dengan pertemanan.					
4	Pemimpin di tempat saya mengajar mampu menjadi teladan dalam menguasai suatu pemasalahan.					
5	Pemimpin di tempat saya mengajar memberikan reward untuk setiap guru yang rajin, ramah, dan mampu mengayomi murid.					

### Disiplin (X3)

No	Pernyataan	SS	S	KS	TS	STS
1	Saya selalu mentaati ketentuan jam kerja, masuk, pulang dan istirahat tepat waktu sesuai peraturan yang berlaku.					
2	Saya membuat RPP untuk semua materi pembelajaran yang akan saya ajarkan.					
3	Saya mengenakan seragam kerja sesuai hari yang telah ditentukan.					
4	Saya selalu hadir dikelas dan mengakhiri kelas dengan tepat waktu.					
5	Saya menata dan meletakkan kembali media pembelajaran pada tempatnya setelah selesai menggunakannya.					

### Kinerja (Y)

No	Pernyataan	SS	S	KS	TS	STS
1	Guru merencanakan kegiatan pembelajaran yang saling terkait satu sama lain serta melaksanakan evaluasi pembelajaran.					
2	Guru menggunakan alat bantu mengajar/audio visual untuk meningkatkan motivasi belajar peserta didik dan mempelajari materi pembelajaran sebelum proses belajar mengajar dimulai.					
3	Guru menciptakan ketertiban, kedisiplinan dan kenyamanan dalam proses belajar serta melakukan penelitian terhadap hasil belajar siswa.					
4	Guru memberikan kesempatan pada peserta didik untuk mengemukakan pendapatnya dan menegembangkan suasana belajar kelompok yang aktif dan kreatif sebagai metode dalam pembelajaran.					
5	Guru menganalisis hasil penilaian dari proses belajar untuk mengetahui dan menentukan ketuntasan siswa dalam pembelajaran.					

## Data Hasil Kuesioner

X1.1	X1.2	X1.3	X1.4	X1.5	Total X1	X2.1	X2.2	X2.3	X2.4	X2.5	Total X2	X3.1	X3.2	X3.3	X3.4	X3.5	Total X3	Y1.1	Y1.2	Y1.3	Y1.4	Y1.5	Total Y1
5	5	4	4	5	23	5	5	5	4	5	24	4	5	5	5	5	24	5	5	5	5	5	25
5	5	4	4	4	22	4	5	4	4	4	21	4	5	5	4	4	22	4	4	4	4	5	21
4	4	4	4	4	20	4	5	4	5	4	22	4	5	4	4	4	21	4	4	4	4	4	20
4	4	5	5	4	22	5	5	4	5	4	23	5	4	5	5	5	24	5	4	4	5	4	22
4	4	4	4	4	20	4	4	4	3	4	19	4	4	4	4	4	20	4	4	4	3	4	19
5	4	4	4	4	21	4	4	4	5	5	22	5	4	4	4	4	21	4	4	4	5	5	22
5	5	5	4	5	24	5	4	5	5	4	23	5	5	5	5	5	25	4	5	5	5	5	24
5	4	4	4	4	21	4	5	5	4	4	22	4	4	4	5	5	22	4	4	4	5	4	21
5	4	5	5	5	24	4	5	5	5	4	23	4	4	4	5	5	22	5	5	4	5	5	24
5	5	4	5	4	23	4	5	5	5	5	24	4	5	3	4	4	20	4	5	4	5	4	22
5	4	5	5	4	23	5	5	5	5	5	25	5	5	5	5	4	24	5	5	5	5	5	25
4	5	5	4	4	22	5	4	5	4	5	23	5	4	5	5	5	24	4	5	5	5	5	24
5	5	5	5	5	25	5	5	4	5	5	24	4	5	4	5	4	22	5	4	5	5	5	24
4	5	5	5	5	24	4	5	4	4	5	22	4	4	5	5	4	22	5	4	5	4	4	22
3	3	3	3	3	15	3	3	3	3	3	15	3	3	3	3	3	15	2	3	3	3	3	14
4	4	4	5	4	21	4	5	4	4	5	22	4	4	3	4	4	19	4	5	4	4	4	21
5	5	4	5	4	23	5	5	4	4	3	21	5	5	5	5	5	25	4	4	4	5	4	21
4	4	4	5	5	22	4	4	4	4	4	20	4	4	4	4	5	21	4	4	5	5	4	22
4	5	4	4	4	21	4	5	4	5	5	23	4	4	4	5	5	22	4	5	5	4	5	23
4	4	4	4	4	20	4	3	4	3	4	18	4	4	4	4	5	21	4	5	4	4	4	21
4	4	4	5	4	21	4	5	4	5	3	21	4	4	4	4	4	20	4	4	4	4	4	20
5	5	4	5	5	24	5	5	4	4	5	23	4	4	5	4	5	22	4	4	5	5	4	22
3	3	3	3	3	15	3	3	3	3	3	15	3	3	3	3	3	15	3	3	3	3	3	15
4	4	5	5	5	23	5	4	5	5	5	24	4	4	4	5	4	21	4	4	5	5	4	22
4	4	5	4	4	21	4	4	4	4	4	20	4	4	4	5	3	20	4	4	4	4	4	20
5	5	5	5	5	25	4	5	5	4	3	21	4	5	5	4	5	23	4	4	5	4	5	22
5	5	5	5	5	25	5	4	5	5	5	24	5	5	4	5	5	24	5	5	5	5	5	25
4	4	4	4	4	20	5	4	5	3	4	21	5	5	5	5	5	25	4	4	5	4	5	22
4	4	4	4	4	20	4	4	4	4	3	19	4	3	3	4	4	18	4	4	4	4	4	20
5	5	5	5	5	25	5	5	5	4	5	24	3	3	5	5	4	20	5	5	5	5	5	25

Tabel r untuk df = 1 - 50

df = (N-2)	Tingkat signifikansi untuk uji satu arah				
	0.05	0.025	0.01	0.005	0.0005
	Tingkat signifikansi untuk uji dua arah				
	0.1	0.05	0.02	0.01	0.001
1	0.9877	0.9969	0.9995	0.9999	1.0000
2	0.9000	0.9500	0.9800	0.9900	0.9990
3	0.8054	0.8783	0.9343	0.9587	0.9911
4	0.7293	0.8114	0.8822	0.9172	0.9741
5	0.6694	0.7545	0.8329	0.8745	0.9509
6	0.6215	0.7067	0.7887	0.8343	0.9249
7	0.5822	0.6664	0.7498	0.7977	0.8983
8	0.5494	0.6319	0.7155	0.7646	0.8721
9	0.5214	0.6021	0.6851	0.7348	0.8470
10	0.4973	0.5760	0.6581	0.7079	0.8233
11	0.4762	0.5529	0.6339	0.6835	0.8010
12	0.4575	0.5324	0.6120	0.6614	0.7800
13	0.4409	0.5140	0.5923	0.6411	0.7604
14	0.4259	0.4973	0.5742	0.6226	0.7419
15	0.4124	0.4821	0.5577	0.6055	0.7247
16	0.4000	0.4683	0.5425	0.5897	0.7084
17	0.3887	0.4555	0.5285	0.5751	0.6932
18	0.3783	0.4438	0.5155	0.5614	0.6788
19	0.3687	0.4329	0.5034	0.5487	0.6652
20	0.3598	0.4227	0.4921	0.5368	0.6524
21	0.3515	0.4132	0.4815	0.5256	0.6402
22	0.3438	0.4044	0.4716	0.5151	0.6287
23	0.3365	0.3961	0.4622	0.5052	0.6178
24	0.3297	0.3882	0.4534	0.4958	0.6074
25	0.3233	0.3809	0.4451	0.4869	0.5974
26	0.3172	0.3739	0.4372	0.4785	0.5880
27	0.3115	0.3673	0.4297	0.4705	0.5790
28	0.3061	0.3610	0.4226	0.4629	0.5703
29	0.3009	0.3550	0.4158	0.4556	0.5620
30	0.2960	0.3494	0.4093	0.4487	0.5541
31	0.2913	0.3440	0.4032	0.4421	0.5465
32	0.2869	0.3388	0.3972	0.4357	0.5392
33	0.2826	0.3338	0.3916	0.4296	0.5322
34	0.2785	0.3291	0.3862	0.4238	0.5254
35	0.2746	0.3246	0.3810	0.4182	0.5189
36	0.2709	0.3202	0.3760	0.4128	0.5126
37	0.2673	0.3160	0.3712	0.4076	0.5066
38	0.2638	0.3120	0.3665	0.4026	0.5007
39	0.2605	0.3081	0.3621	0.3978	0.4950
40	0.2573	0.3044	0.3578	0.3932	0.4896
41	0.2542	0.3008	0.3536	0.3887	0.4843
42	0.2512	0.2973	0.3496	0.3843	0.4791
43	0.2483	0.2940	0.3457	0.3801	0.4742
44	0.2455	0.2907	0.3420	0.3761	0.4694
45	0.2429	0.2876	0.3384	0.3721	0.4647
46	0.2403	0.2845	0.3348	0.3683	0.4601
47	0.2377	0.2816	0.3314	0.3646	0.4557
48	0.2353	0.2787	0.3281	0.3610	0.4514
49	0.2329	0.2759	0.3249	0.3575	0.4473
50	0.2306	0.2732	0.3218	0.3542	0.4432

**Titik Persentase Distribusi t (df = 1 – 40)**

	Pr	0.25	0.10	0.05	0.025	0.01	0.005	0.001
df		0.50	0.20	0.10	0.050	0.02	0.010	0.002
1	1.00000	3.07768	6.31375	12.70620	31.82052	63.65674	318.30884	
2	0.81650	1.88562	2.91999	4.30265	6.96456	9.92484	22.32712	
3	0.76489	1.63774	2.35336	3.18245	4.54070	5.84091	10.21453	
4	0.74070	1.53321	2.13185	2.77645	3.74695	4.60409	7.17318	
5	0.72669	1.47588	2.01505	2.57058	3.36493	4.03214	5.89343	
6	0.71756	1.43976	1.94318	2.44691	3.14267	3.70743	5.20763	
7	0.71114	1.41492	1.89458	2.36462	2.99795	3.49948	4.78529	
8	0.70639	1.39682	1.85955	2.30600	2.89646	3.35539	4.50079	
9	0.70272	1.38303	1.83311	2.26216	2.82144	3.24984	4.29681	
10	0.69981	1.37218	1.81246	2.22814	2.76377	3.16927	4.14370	
11	0.69745	1.36343	1.79588	2.20099	2.71808	3.10581	4.02470	
12	0.69548	1.35622	1.78229	2.17881	2.68100	3.05454	3.92963	
13	0.69383	1.35017	1.77093	2.16037	2.65031	3.01228	3.85198	
14	0.69242	1.34503	1.76131	2.14479	2.62449	2.97684	3.78739	
15	0.69120	1.34061	1.75305	2.13145	2.60248	2.94671	3.73283	
16	0.69013	1.33676	1.74588	2.11991	2.58349	2.92078	3.68615	
17	0.68920	1.33338	1.73961	2.10982	2.56693	2.89823	3.64577	
18	0.68836	1.33039	1.73406	2.10092	2.55238	2.87844	3.61048	
19	0.68762	1.32773	1.72913	2.09302	2.53948	2.86093	3.57940	
20	0.68695	1.32534	1.72472	2.08596	2.52798	2.84534	3.55181	
21	0.68635	1.32319	1.72074	2.07961	2.51765	2.83136	3.52715	
22	0.68581	1.32124	1.71714	2.07387	2.50832	2.81876	3.50499	
23	0.68531	1.31946	1.71387	2.06866	2.49987	2.80734	3.48496	
24	0.68485	1.31784	1.71088	2.06390	2.49216	2.79694	3.46678	
25	0.68443	1.31635	1.70814	2.05954	2.48511	2.78744	3.45019	
26	0.68404	1.31497	1.70562	2.05553	2.47863	2.77871	3.43500	
27	0.68368	1.31370	1.70329	2.05183	2.47266	2.77068	3.42103	
28	0.68335	1.31253	1.70113	2.04841	2.46714	2.76326	3.40816	
29	0.68304	1.31143	1.69913	2.04523	2.46202	2.75639	3.39624	
30	0.68276	1.31042	1.69726	2.04227	2.45726	2.75000	3.38518	
31	0.68249	1.30946	1.69552	2.03951	2.45282	2.74404	3.37490	
32	0.68223	1.30857	1.69389	2.03693	2.44868	2.73848	3.36531	
33	0.68200	1.30774	1.69236	2.03452	2.44479	2.73328	3.35634	
34	0.68177	1.30695	1.69092	2.03224	2.44115	2.72839	3.34793	
35	0.68156	1.30621	1.68957	2.03011	2.43772	2.72381	3.34005	
36	0.68137	1.30551	1.68830	2.02809	2.43449	2.71948	3.33262	
37	0.68118	1.30485	1.68709	2.02619	2.43145	2.71541	3.32563	
38	0.68100	1.30423	1.68595	2.02439	2.42857	2.71156	3.31903	
39	0.68083	1.30364	1.68488	2.02269	2.42584	2.70791	3.31279	
40	0.68067	1.30308	1.68385	2.02108	2.42326	2.70446	3.30688	



**Titik Persentase Distribusi F untuk Probabilita = 0,05**

df untuk penyebut (N2)	df untuk pembilang (N1)														
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
1	161	199	216	225	230	234	237	239	241	242	243	244	245	245	246
2	18.51	19.00	19.16	19.25	19.30	19.33	19.35	19.37	19.38	19.40	19.40	19.41	19.42	19.42	19.43
3	10.13	9.55	9.28	9.12	9.01	8.94	8.89	8.85	8.81	8.79	8.76	8.74	8.73	8.71	8.70
4	7.71	6.94	6.59	6.39	6.26	6.16	6.09	6.04	6.00	5.96	5.94	5.91	5.89	5.87	5.86
5	6.61	5.79	5.41	5.19	5.05	4.95	4.88	4.82	4.77	4.74	4.70	4.68	4.66	4.64	4.62
6	5.99	5.14	4.76	4.53	4.39	4.28	4.21	4.15	4.10	4.06	4.03	4.00	3.98	3.96	3.94
7	5.59	4.74	4.35	4.12	3.97	3.87	3.79	3.73	3.68	3.64	3.60	3.57	3.55	3.53	3.51
8	5.32	4.46	4.07	3.84	3.69	3.58	3.50	3.44	3.39	3.35	3.31	3.28	3.26	3.24	3.22
9	5.12	4.26	3.86	3.63	3.48	3.37	3.29	3.23	3.18	3.14	3.10	3.07	3.05	3.03	3.01
10	4.96	4.10	3.71	3.48	3.33	3.22	3.14	3.07	3.02	2.98	2.94	2.91	2.89	2.86	2.85
11	4.84	3.98	3.59	3.36	3.20	3.09	3.01	2.95	2.90	2.85	2.82	2.79	2.76	2.74	2.72
12	4.75	3.89	3.49	3.26	3.11	3.00	2.91	2.85	2.80	2.75	2.72	2.69	2.66	2.64	2.62
13	4.67	3.81	3.41	3.18	3.03	2.92	2.83	2.77	2.71	2.67	2.63	2.60	2.58	2.55	2.53
14	4.60	3.74	3.34	3.11	2.96	2.85	2.76	2.70	2.65	2.60	2.57	2.53	2.51	2.48	2.46
15	4.54	3.68	3.29	3.06	2.90	2.79	2.71	2.64	2.59	2.54	2.51	2.48	2.45	2.42	2.40
16	4.49	3.63	3.24	3.01	2.85	2.74	2.66	2.59	2.54	2.49	2.46	2.42	2.40	2.37	2.35
17	4.45	3.59	3.20	2.96	2.81	2.70	2.61	2.55	2.49	2.45	2.41	2.38	2.35	2.33	2.31
18	4.41	3.55	3.16	2.93	2.77	2.66	2.58	2.51	2.46	2.41	2.37	2.34	2.31	2.29	2.27
19	4.38	3.52	3.13	2.90	2.74	2.63	2.54	2.48	2.42	2.38	2.34	2.31	2.28	2.26	2.23
20	4.35	3.49	3.10	2.87	2.71	2.60	2.51	2.45	2.39	2.35	2.31	2.28	2.25	2.22	2.20
21	4.32	3.47	3.07	2.84	2.68	2.57	2.49	2.42	2.37	2.32	2.28	2.25	2.22	2.20	2.18
22	4.30	3.44	3.05	2.82	2.66	2.55	2.46	2.40	2.34	2.30	2.26	2.23	2.20	2.17	2.15
23	4.28	3.42	3.03	2.80	2.64	2.53	2.44	2.37	2.32	2.27	2.24	2.20	2.18	2.15	2.13
24	4.26	3.40	3.01	2.78	2.62	2.51	2.42	2.36	2.30	2.25	2.22	2.18	2.15	2.13	2.11
25	4.24	3.39	2.99	2.76	2.60	2.49	2.40	2.34	2.28	2.24	2.20	2.16	2.14	2.11	2.09
26	4.23	3.37	2.98	2.74	2.59	2.47	2.39	2.32	2.27	2.22	2.18	2.15	2.12	2.09	2.07
27	4.21	3.35	2.96	2.73	2.57	2.46	2.37	2.31	2.25	2.20	2.17	2.13	2.10	2.08	2.06
28	4.20	3.34	2.95	2.71	2.56	2.45	2.36	2.29	2.24	2.19	2.15	2.12	2.09	2.06	2.04
29	4.18	3.33	2.93	2.70	2.55	2.43	2.35	2.28	2.22	2.18	2.14	2.10	2.08	2.05	2.03
30	4.17	3.32	2.92	2.69	2.53	2.42	2.33	2.27	2.21	2.16	2.13	2.09	2.06	2.04	2.01
31	4.16	3.30	2.91	2.68	2.52	2.41	2.32	2.25	2.20	2.15	2.11	2.08	2.05	2.03	2.00
32	4.15	3.29	2.90	2.67	2.51	2.40	2.31	2.24	2.19	2.14	2.10	2.07	2.04	2.01	1.99
33	4.14	3.28	2.89	2.66	2.50	2.39	2.30	2.23	2.18	2.13	2.09	2.06	2.03	2.00	1.98
34	4.13	3.28	2.88	2.65	2.49	2.38	2.29	2.23	2.17	2.12	2.08	2.05	2.02	1.99	1.97
35	4.12	3.27	2.87	2.64	2.49	2.37	2.29	2.22	2.16	2.11	2.07	2.04	2.01	1.99	1.96
36	4.11	3.26	2.87	2.63	2.48	2.36	2.28	2.21	2.15	2.11	2.07	2.03	2.00	1.98	1.95
37	4.11	3.25	2.86	2.63	2.47	2.36	2.27	2.20	2.14	2.10	2.06	2.02	2.00	1.97	1.95
38	4.10	3.24	2.85	2.62	2.46	2.35	2.26	2.19	2.14	2.09	2.05	2.02	1.99	1.96	1.94
39	4.09	3.24	2.85	2.61	2.46	2.34	2.26	2.19	2.13	2.08	2.04	2.01	1.98	1.95	1.93
40	4.08	3.23	2.84	2.61	2.45	2.34	2.25	2.18	2.12	2.08	2.04	2.00	1.97	1.95	1.92
41	4.08	3.23	2.83	2.60	2.44	2.33	2.24	2.17	2.12	2.07	2.03	2.00	1.97	1.94	1.92
42	4.07	3.22	2.83	2.59	2.44	2.32	2.24	2.17	2.11	2.06	2.03	1.99	1.96	1.94	1.91
43	4.07	3.21	2.82	2.59	2.43	2.32	2.23	2.16	2.11	2.06	2.02	1.99	1.96	1.93	1.91
44	4.06	3.21	2.82	2.58	2.43	2.31	2.23	2.16	2.10	2.05	2.01	1.98	1.95	1.92	1.90
45	4.06	3.20	2.81	2.58	2.42	2.31	2.22	2.15	2.10	2.05	2.01	1.97	1.94	1.92	1.89

## Dokumentasi





Your temporary usage period for IBM SPSS Statistics will expire in 4678 days.

```
CORRELATIONS
/VARIABLES=X1.1 X1.2 X1.3 X1.4 X1.5 TOTAL
/PRINT=TWOTAIL NOSIG
/MISSING=PAIRWISE.
```

## Correlations

<b>Notes</b>		
Output Created		11-MAR-2023 15:51:17
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax		CORRELATIONS /VARIABLES=X1.1 X1.2 X1.3 X1.4 X1.5 TOTAL /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.11

[DataSet0]

		<b>Correlations</b>					
		X1.1	X1.2	X1.3	X1.4	X1.5	TOTAL
X1.1	Pearson Correlation	1	.686**	.457*	.514**	.596**	.801**
	Sig. (2-tailed)		.000	.011	.004	.001	.000
	N	30	30	30	30	30	30
X1.2	Pearson Correlation	.686**	1	.493**	.469**	.630**	.806**
	Sig. (2-tailed)	.000		.006	.009	.000	.000
	N	30	30	30	30	30	30

X1.3	Pearson Correlation	.457*	.493**	1	.605**	.668**	.791**
	Sig. (2-tailed)	.011	.006		.000	.000	.000
	N	30	30	30	30	30	30
X1.4	Pearson Correlation	.514**	.469**	.605**	1	.656**	.799**
	Sig. (2-tailed)	.004	.009	.000		.000	.000
	N	30	30	30	30	30	30
X1.5	Pearson Correlation	.596**	.630**	.668**	.656**	1	.871**
	Sig. (2-tailed)	.001	.000	.000	.000		.000
	N	30	30	30	30	30	30
TOTAL	Pearson Correlation	.801**	.806**	.791**	.799**	.871**	1
	Sig. (2-tailed)	.000	.000	.000	.000	.000	
	N	30	30	30	30	30	30

\*\* . Correlation is significant at the 0.01 level (2-tailed).

\* . Correlation is significant at the 0.05 level (2-tailed).

#### RELIABILITY

```

/VARIABLES=X1.1 X1.2 X1.3 X1.4 X1.5
/SCALE('ALL VARIABLES') ALL
/MODEL=ALPHA
/SUMMARY=TOTAL.

```

## Reliability

### Notes

Output Created	11-MAR-2023 15:51:44	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
	Matrix Input	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure.

Syntax	RELIABILITY /VARIABLES=X1.1 X1.2 X1.3 X1.4 X1.5 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA /SUMMARY=TOTAL.
Resources	Processor Time 00:00:00.03 Elapsed Time 00:00:00.01

## Scale: ALL VARIABLES

### Case Processing Summary

		N	%
Cases	Valid	30	100.0
	Excluded <sup>a</sup>	0	.0
	Total	30	100.0

a. Listwise deletion based on all variables in the procedure.

### Reliability Statistics

Cronbach's Alpha	N of Items
.872	5

### Item-Total Statistics

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Cronbach's Alpha if Item Deleted
X1.1	17.43	4.116	.678	.850
X1.2	17.47	4.120	.687	.847
X1.3	17.50	4.190	.667	.852
X1.4	17.40	4.110	.674	.851
X1.5	17.53	3.982	.789	.823

### CORRELATIONS

```

/VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5 TOTAL
/PRINT=TWOTAIL NOSIG
/MISSING=PAIRWISE.

```

## Correlations

<b>Notes</b>		
Output Created		11-MAR-2023 15:53:36
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax		CORRELATIONS /VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5 TOTAL /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.02

## Correlations

		X2.1	X2.2	X2.3	X2.4	X2.5	TOTAL
X2.1	Pearson Correlation	1	.362*	.625**	.365*	.490**	.755**
	Sig. (2-tailed)		.050	.000	.048	.006	.000
	N	30	30	30	30	30	30
X2.2	Pearson Correlation	.362*	1	.362*	.538**	.310	.704**
	Sig. (2-tailed)	.050		.050	.002	.096	.000
	N	30	30	30	30	30	30
X2.3	Pearson Correlation	.625**	.362*	1	.365*	.416*	.732**
	Sig. (2-tailed)	.000	.050		.048	.022	.000
	N	30	30	30	30	30	30
X2.4	Pearson Correlation	.365*	.538**	.365*	1	.390*	.739**
	Sig. (2-tailed)	.048	.002	.048		.033	.000
	N	30	30	30	30	30	30

X2.5	Pearson Correlation	.490**	.310	.416*	.390*	1	.731**
	Sig. (2-tailed)	.006	.096	.022	.033		.000
	N	30	30	30	30	30	30
TOTAL	Pearson Correlation	.755**	.704**	.732**	.739**	.731**	1
	Sig. (2-tailed)	.000	.000	.000	.000	.000	
	N	30	30	30	30	30	30

\*. Correlation is significant at the 0.05 level (2-tailed).

\*\* Correlation is significant at the 0.01 level (2-tailed).

#### RELIABILITY

```

/VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5
/SCALE('ALL VARIABLES') ALL
/MODEL=ALPHA
/SUMMARY=TOTAL.

```

## Reliability

### Notes

Output Created	11-MAR-2023 15:53:48	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
	Matrix Input	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure.
Syntax	RELIABILITY /VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA /SUMMARY=TOTAL.	
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.01



## Scale: ALL VARIABLES

### Case Processing Summary

		N	%
Cases	Valid	30	100.0
	Excluded <sup>a</sup>	0	.0
	Total	30	100.0

a. Listwise deletion based on all variables in the procedure.

### Reliability Statistics

Cronbach's Alpha	N of Items
.779	5

### Item-Total Statistics

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Cronbach's Alpha if Item Deleted
X2.1	17.27	4.271	.614	.721
X2.2	17.13	4.257	.518	.749
X2.3	17.27	4.340	.582	.731
X2.4	17.37	4.033	.553	.739
X2.5	17.37	3.964	.524	.752

### CORRELATIONS

```

/VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5 TOTAL
/PRINT=TWOTAIL NOSIG
/MISSING=PAIRWISE.

```

## Correlations

### Notes

Output Created	11-MAR-2023 15:54:22	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30

Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax		CORRELATIONS /VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5 TOTAL /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.
Resources	Processor Time	00:00:00.03
	Elapsed Time	00:00:00.07

```
RELIABILITY
/VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5
/SCALE('ALL VARIABLES') ALL
/MODEL=ALPHA
/SUMMARY=TOTAL.
```

## Reliability

Notes		
Output Created		11-MAR-2023 15:54:37
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
	Matrix Input	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure.

Syntax	RELIABILITY /VARIABLES=X2.1 X2.2 X2.3 X2.4 X2.5 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA /SUMMARY=TOTAL.	
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Scale: ALL VARIABLES

### CORRELATIONS

```

/VARIABLES=X3.1 X3.2 X3.3 X3.4 X3.5 TOTAL
/PRINT=TWOTAIL NOSIG
/MISSING=PAIRWISE.

```

## Correlations

Notes		
Output Created	11-MAR-2023 15:56:38	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax	CORRELATIONS /VARIABLES=X3.1 X3.2 X3.3 X3.4 X3.5 TOTAL /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.	
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.01

### Correlations

		X3.1	X3.2	X3.3	X3.4	X3.5	TOTAL
X3.1	Pearson Correlation	1	.586**	.467**	.525**	.537**	.794**
	Sig. (2-tailed)		.001	.009	.003	.002	.000
	N	30	30	30	30	30	30
X3.2	Pearson Correlation	.586**	1	.444*	.382*	.413*	.734**
	Sig. (2-tailed)	.001		.014	.037	.023	.000
	N	30	30	30	30	30	30
X3.3	Pearson Correlation	.467**	.444*	1	.583**	.527**	.796**
	Sig. (2-tailed)	.009	.014		.001	.003	.000
	N	30	30	30	30	30	30
X3.4	Pearson Correlation	.525**	.382*	.583**	1	.481**	.765**
	Sig. (2-tailed)	.003	.037	.001		.007	.000
	N	30	30	30	30	30	30
X3.5	Pearson Correlation	.537**	.413*	.527**	.481**	1	.768**
	Sig. (2-tailed)	.002	.023	.003	.007		.000
	N	30	30	30	30	30	30
TOTAL	Pearson Correlation	.794**	.734**	.796**	.765**	.768**	1
	Sig. (2-tailed)	.000	.000	.000	.000	.000	
	N	30	30	30	30	30	30

\*\* . Correlation is significant at the 0.01 level (2-tailed).

\* . Correlation is significant at the 0.05 level (2-tailed).

#### RELIABILITY

```

/VARIABLES=X3.1 X3.2 X3.3 X3.4 X3.5
/SCALE('ALL VARIABLES') ALL
/MODEL=ALPHA
/SUMMARY=TOTAL.

```

## Reliability

### Notes

Output Created	11-MAR-2023 15:56:48	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>

	<u>N of Rows in Working Data File</u>	30
	Matrix Input	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure.
Syntax		RELIABILITY /VARIABLES=X3.1 X3.2 X3.3 X3.4 X3.5 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA /SUMMARY=TOTAL.
Resources	<u>Processor Time</u>	00:00:00.00
	Elapsed Time	00:00:00.01

## Scale: ALL VARIABLES

### Case Processing Summary

		N	%
Cases	Valid	30	100.0
	Excluded <sup>a</sup>	0	.0
	Total	30	100.0

a. Listwise deletion based on all variables in the procedure.

### Reliability Statistics

Cronbach's Alpha	N of Items
.828	5

### Item-Total Statistics

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Cronbach's Alpha if Item Deleted
X3.1	17.30	4.424	.678	.781
X3.2	17.23	4.392	.566	.810
X3.3	17.23	4.047	.644	.788
X3.4	17.00	4.414	.626	.793

X3.5	17.10	4.300	.619	.795
------	-------	-------	------	------

```

CORRELATIONS
/VARIABLES=Y1 Y2 Y3 Y4 Y5 TOTAL
/PRINT=TWOTAIL NOSIG
/MISSING=PAIRWISE.

```

## Correlations

		Notes
Output Created		11-MAR-2023 16:02:52
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each pair of variables are based on all the cases with valid data for that pair.
Syntax		CORRELATIONS /VARIABLES=Y1 Y2 Y3 Y4 Y5 TOTAL /PRINT=TWOTAIL NOSIG /MISSING=PAIRWISE.
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.01

## Correlations

		Y1	Y2	Y3	Y4	Y5	TOTAL
Y1	Pearson Correlation	1	.581**	.600**	.614**	.620**	.840**
	Sig. (2-tailed)		.001	.000	.000	.000	.000
	N	30	30	30	30	30	30
Y2	Pearson Correlation	.581**	1	.503**	.520**	.630**	.786**
	Sig. (2-tailed)	.001		.005	.003	.000	.000

	N	30	30	30	30	30	30
Y3	Pearson Correlation	.600**	.503**	1	.556**	.686**	.818**
	Sig. (2-tailed)	.000	.005		.001	.000	.000
	N	30	30	30	30	30	30
Y4	Pearson Correlation	.614**	.520**	.556**	1	.515**	.795**
	Sig. (2-tailed)	.000	.003	.001		.004	.000
	N	30	30	30	30	30	30
Y5	Pearson Correlation	.620**	.630**	.686**	.515**	1	.841**
	Sig. (2-tailed)	.000	.000	.000	.004		.000
	N	30	30	30	30	30	30
TOTAL	Pearson Correlation	.840**	.786**	.818**	.795**	.841**	1
	Sig. (2-tailed)	.000	.000	.000	.000	.000	
	N	30	30	30	30	30	30

\*\* . Correlation is significant at the 0.01 level (2-tailed).

#### RELIABILITY

```

/VARIABLES=Y1 Y2 Y3 Y4 Y5
/SCALE ('ALL VARIABLES') ALL
/MODEL=ALPHA
/SUMMARY=TOTAL.

```

## Reliability

### Notes

Output Created	11-MAR-2023 16:03:05	
Comments		
Input	Active Dataset	DataSet0
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	30
	Matrix Input	
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics are based on all cases with valid data for all variables in the procedure.

Syntax		RELIABILITY /VARIABLES=Y1 Y2 Y3 Y4 Y5 /SCALE('ALL VARIABLES') ALL /MODEL=ALPHA /SUMMARY=TOTAL.
Resources	Processor Time	00:00:00.00
	Elapsed Time	00:00:00.00

## Scale: ALL VARIABLES

### Case Processing Summary

		N	%
Cases	Valid	30	100.0
	Excluded <sup>a</sup>	0	.0
	Total	30	100.0

a. Listwise deletion based on all variables in the procedure.

### Reliability Statistics

Cronbach's Alpha	N of Items
.874	5

### Item-Total Statistics

	Scale Mean if Item Deleted	Scale Variance if Item Deleted	Corrected Item-Total Correlation	Cronbach's Alpha if Item Deleted
Y1	17.50	4.259	.735	.839
Y2	17.37	4.585	.667	.855
Y3	17.27	4.409	.708	.845
Y4	17.23	4.323	.659	.859
Y5	17.30	4.355	.744	.837



### Variables Entered/Removed<sup>a</sup>

Model	Variables Entered	Variables Removed	Method
1	Disiplin (X3), Motivasi (X2), Kepemimpinan (X1) <sup>b</sup>		Enter

a. Dependent Variable: Kinerja (Y)

b. All requested variables entered.

### Model Summary<sup>b</sup>

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.935 <sup>a</sup>	.874	.859	.966

a. Predictors: (Constant), Disiplin (X3), Motivasi (X2), Kepemimpinan (X1)

b. Dependent Variable: Kinerja (Y)

### ANOVA<sup>a</sup>

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	168.392	3	56.131	60.121	.000 <sup>b</sup>
	Residual	24.274	26	.934		
	Total	192.667	29			

a. Dependent Variable: Kinerja (Y)

b. Predictors: (Constant), Disiplin (X3), Motivasi (X2), Kepemimpinan (X1)

### Coefficients<sup>a</sup>

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.	Collinearity Statistics	
		B	Std. Error	Beta			Tolerance	VIF
1	(Constant)	-1.073	1.717		-.625	.538		
	Kepemimpinan (X1)	.323	.135	.313	2.400	.024	.286	3.502
	Motivasi (X2)	.494	.135	.476	3.666	.001	.287	3.482
	Disiplin (X3)	.234	.101	.231	2.306	.029	.485	2.062

a. Dependent Variable: Kinerja (Y)

### Collinearity Diagnostics<sup>a</sup>

Model	Dimension	Eigenvalue	Condition Index	(Constant)	Variance Proportions		
					Kepemimpinan (X1)	Motivasi (X2)	Disiplin (X3)
1	1	3.985	1.000	.00	.00	.00	.00
	2	.008	22.405	1.00	.04	.05	.05
	3	.005	29.103	.00	.11	.12	.95
	4	.002	43.143	.00	.85	.83	.00

a. Dependent Variable: Kinerja (Y)

### Residuals Statistics<sup>a</sup>


	Minimum	Maximum	Mean	Std. Deviation	N
Predicted Value	14.69	24.47	21.67	2.410	30
Std. Predicted Value	-2.896	1.162	.000	1.000	30
Standard Error of Predicted Value	.208	.553	.339	.098	30
Adjusted Predicted Value	14.54	24.41	21.67	2.407	30
Residual	-1.573	1.813	.000	.915	30
Std. Residual	-1.628	1.876	.000	.947	30
Stud. Residual	-1.826	2.044	.000	1.024	30
Deleted Residual	-1.979	2.152	.000	1.074	30
Stud. Deleted Residual	-1.918	2.188	.004	1.046	30
Mahal. Distance	.384	8.536	2.900	2.352	30
Cook's Distance	.000	.228	.045	.061	30
Centered Leverage Value	.013	.294	.100	.081	30

a. Dependent Variable: Kinerja (Y)

# Charts

Normal P-P Plot of Regression Standardized Residual

Dependent Variable: Kinerja (Y)



Scatterplot

Dependent Variable: Kinerja (Y)

